

Nonrepetitive Colourings of Planar Graphs with $O(\log n)$ Colours

Vida Dujmović[†] Fabrizio Frati[‡] Gwenaël Joret[§] David R. Wood[¶]

February 8, 2012

Abstract

A vertex colouring of a graph is *nonrepetitive* if there is no path for which the first half of the path is assigned the same sequence of colours as the second half. The *nonrepetitive chromatic number* of a graph G is the minimum integer k such that G has a nonrepetitive k -colouring. Whether planar graphs have bounded nonrepetitive chromatic number is one of the most important open problems in the field. Despite this, the best known upper bound is $O(\sqrt{n})$ for n -vertex planar graphs. We prove a $O(\log n)$ upper bound.

1 Introduction

A vertex colouring of a graph is *nonrepetitive* if there is no path for which the first half of the path is assigned the same sequence of colours as the second half. More precisely, a *k -colouring* of a graph G is a function ψ that assigns one of k colours to each vertex of G . A path $(v_1, v_2, \dots, v_{2t})$ of even order in G is *repetitively coloured* by ψ if $\psi(v_i) = \psi(v_{t+i})$ for all $i \in [1, t] := \{1, 2, \dots, t\}$. A colouring ψ of G is *nonrepetitive* if no path of G is repetitively coloured by ψ . Observe that a nonrepetitive colouring is *proper*, in the sense that adjacent vertices are coloured differently. The *nonrepetitive chromatic number* $\pi(G)$ is the minimum integer k such that G admits a nonrepetitive k -colouring.

[†]School of Computer Science, Carleton University, Ottawa, Canada (vida@scs.carleton.ca). Supported by NSERC and an Endeavour Fellowship from the Australian Government.

[‡]School of Information Technologies, The University of Sydney, Sydney, Australia (brillo@it.usyd.edu.au).

[§]Département d’Informatique, Université Libre de Bruxelles, Brussels, Belgium (gjoret@ulb.ac.be). Postdoctoral Researcher of the Fonds National de la Recherche Scientifique (F.R.S.-FNRS). Supported by an Endeavour Fellowship from the Australian Government.

[¶]Department of Mathematics and Statistics, The University of Melbourne, Melbourne, Australia (woodd@unimelb.edu.au). Supported by a QEII Fellowship and a Discovery Project from the Australian Research Council.

The seminal result in this field is by Thue [36], who in 1906 proved that every path is nonrepetitively 3-colourable. Nonrepetitive colourings have recently been widely studied [2–8, 10–12, 18, 20, 22, 24–31, 33–35]; see the surveys [9, 21–23]. A number of graph classes are known to have bounded nonrepetitive chromatic number. In particular, trees are nonrepetitively 4-colourable [7, 29], outerplanar graphs are nonrepetitively 12-colourable [4, 29], and more generally, every graph with treewidth k is nonrepetitively 4^k -colourable [29]. Graphs with maximum degree Δ are nonrepetitively $O(\Delta^2)$ -colourable [3, 21, 22, 26].

Perhaps the most important open problem in the field of nonrepetitive colourings is whether planar graphs have bounded nonrepetitive chromatic number. This question, first asked by Alon et al. [3], has since been mentioned by numerous authors [2, 4, 12, 20–23, 25, 26, 29, 31, 33]. It is widely known that $\pi(G) \in O(\sqrt{n})$ for n -vertex planar graphs¹, and this is the best known upper bound. The best known lower bound is due to Barát and Varjú [4], who constructed a planar graph with nonrepetitive chromatic number at least 10. Here we prove a logarithmic upper bound.

Theorem 1. *For every planar graph G with n vertices,*

$$\pi(G) \leq 8(1 + \log_{3/2} n) .$$

As a secondary contribution, we solve the above open problem when restricted to paths of bounded length.

Theorem 2. *There is a constant c such that, for every integer $k \geq 1$, every planar graph G is c^{k^2} -colourable such that G contains no repetitively coloured path of order at most $2k$.*

Note that the case $k = 2$ of Theorem 2 corresponds to so-called *star colourings*; that is, proper colourings with no 2-coloured P_4 ; see [1, 17, 32, 37]. Albertson et al. [1] proved that every planar graph is star colourable with 20 colours.

2 Proof of Theorem 1

A *layering* of a graph G is a partition V_0, V_1, \dots, V_p of $V(G)$ such that for every edge $vw \in E(G)$, if $v \in V_i$ and $w \in V_j$ then $|i - j| \leq 1$. Each set V_i is called a *layer*. The following lemma by Kündgen and Pelsmajer [29] will be useful.

Lemma 3 ([29]). *For every layering of a graph G , there is a (not necessarily proper) 4-colouring of G such that for every repetitively coloured path $(v_1, v_2, \dots, v_{2t})$, the subpaths (v_1, v_2, \dots, v_t) and $(v_{t+1}, v_{t+2}, \dots, v_{2t})$ have the same layer pattern.*

¹One can prove this bound using a naive application of the Lipton-Tarjan planar separator theorem.

A *separation* of a graph G is a pair (G_1, G_2) of subgraphs of G , such that $G = G_1 \cup G_2$. In particular, there is no edge of G between $V(G_1) - V(G_2)$ and $V(G_2) - V(G_1)$.

Lemma 4. Fix $\epsilon \in (0, 1)$ and $c \geq 1$. Let G be a graph with n vertices. Fix a layering V_0, V_1, \dots, V_p of G . Assume that, for every set $B \subseteq V(G)$, there is a separation (G_1, G_2) of G such that:

- each layer V_i contains at most c vertices in $V(G_1) \cap V(G_2) \cap B$, and
- both $V(G_1) - V(G_2)$ and $V(G_2) - V(G_1)$ contain at most $(1 - \epsilon)|B|$ vertices in B .

Then $\pi(G) \leq 4c(1 + \log_{1/(1-\epsilon)} n)$.

Proof. Run the following recursive algorithm COMPUTE($V(G), 1$).

COMPUTE(B, d)

1. If $B = \emptyset$ then exit.
2. Let (G_1, G_2) be a separation of G such that each layer V_i contains at most c vertices in $V(G_1) \cap V(G_2) \cap B$, and both $V(G_1) - V(G_2)$ and $V(G_2) - V(G_1)$ contain at most $(1 - \epsilon)|B|$ vertices in B .
3. Let $\text{depth}(v) := d$ for each vertex $v \in V(G_1) \cap V(G_2) \cap B$.
4. For $i \in [1, p]$, injectively label the vertices in $V_i \cap V(G_1) \cap V(G_2) \cap B$ by $1, 2, \dots, c$. Let $\text{label}(v)$ be the label assigned to each vertex $v \in V_i \cap V(G_1) \cap V(G_2) \cap B$.
5. COMPUTE($((V(G_1) - V(G_2)) \cap B, d + 1)$
6. COMPUTE($((V(G_2) - V(G_1)) \cap B, d + 1)$

The recursive application of COMPUTE determines a rooted binary tree T , where each node of T corresponds to one call to COMPUTE. Associate each vertex whose depth and label is computed in a particular call to COMPUTE with the corresponding node of T . (Observe that the depth and label of each vertex is determined exactly once.)

Colour each vertex v by $(\text{col}(v), \text{depth}(v), \text{label}(v))$, where col is the 4-colouring from Lemma 3. Suppose on the contrary that $(v_1, v_2, \dots, v_{2t})$ is a repetitively coloured path in G . By Lemma 3, (v_1, v_2, \dots, v_t) and $(v_{t+1}, v_{t+2}, \dots, v_{2t})$ have the same layer pattern. In addition, $\text{depth}(v_i) = \text{depth}(v_{t+i})$ and $\text{label}(v_i) = \text{label}(v_{t+i})$ for all $i \in [1, t]$. Let v_i and v_{t+i} be vertices in this path with minimum depth. Since v_i and v_{t+i} are in the same layer and have the same label, these two vertices were not labelled at the same step of the algorithm. Let x and y be the two nodes of T respectively associated with v_i and v_{t+i} . Let z be the least common ancestor of x and y in T . Say node z corresponds to call COMPUTE(B, d). Thus v_i

and v_{t+i} are in B (since if a vertex v is in B in the call to COMPUTE associated with some node q of T , then v is in B in the call to COMPUTE associated with each ancestor of q in T). Let (G_1, G_2) be the separation in $\text{COMPUTE}(B, d)$. Since $\text{depth}(v_i) = \text{depth}(v_{t+i}) > d$, neither v_i nor v_{t+i} are in $V(G_1) \cap V(G_2)$. Since z is the least common ancestor of x and y , without loss of generality, $v_i \in V(G_1) - V(G_2)$ and $v_{t+i} \in V(G_2) - V(G_1)$. Thus some vertex v_j in the subpath $(v_{i+1}, v_{i+2}, \dots, v_{t+i-1})$ is in $V(G_1) \cap V(G_2)$. If $v_j \in B$ then $\text{depth}(v_j) = d$. If $v_j \notin B$ then $\text{depth}(v_j) < d$. In both cases, $\text{depth}(v_j) < \text{depth}(v_i) = \text{depth}(v_{t+i})$, which contradicts the choice of v_i and v_{t+i} . Hence there is no repetitively coloured path in G .

Observe that the maximum depth is at most $1 + \log_{1/(1-\epsilon)} n$. Therefore the number of colours is at most $4c(1 + \log_{1/(1-\epsilon)} n)$. \square

We now prove that the condition in Lemma 4 holds for *plane triangulations*; that is, embedded planar graphs in which every face is a triangle. If r is a vertex of a connected graph G and V_i is the set of vertices in G at distance i from r , then V_0, V_1, V_2, \dots is a layering of G , called the *layering starting at r* . Observe that for each vertex $v \in V_i$ there is a vr -path that contains exactly one vertex from each layer V_0, V_1, \dots, V_i ; we call this a *monotone* path.

Lemma 5. *Let r be a vertex in a plane triangulation G . Let V_0, V_1, \dots, V_p be the layering of G starting at r . For every set $B \subseteq V(G)$, there is a separation (G_1, G_2) of G such that:*

- *each layer V_i contains at most two vertices in $V(G_1) \cap V(G_2) \cap B$,*
- *both $V(G_1) - V(G_2)$ and $V(G_2) - V(G_1)$ contain at most $\frac{2}{3}|B|$ vertices in B .*

Proof. If $|B| \leq 2$ then $G_1 := G_2 := G$ satisfy the claim. Now assume that $|B| \geq 3$. A *lollipop* S of height k is a walk in G such that:

- either $S = (u_0, u_1, \dots, u_{k-1}, u_k, v_k, v_{k-1}, \dots, v_1, v_0)$ as in Figure 1(a),
or $S = (u_0, u_1, \dots, u_{k-1}, u_k, u_{k+1}, v_k, v_{k-1}, \dots, v_1, v_0)$ as in Figure 1(b),
where $u_i, v_i \in V_i$ for each $i \in [1, k]$, and $u_{k+1} \in V_{k+1}$;
- $u_0 = v_0 = r$ and $u_k \neq v_k$; and
- if $u_i = v_i$ for some $i \in [1, k-1]$, then $u_j = v_j$ for each $j \in [0, i]$.

Consider a lollipop S . We define vertices to the right and left of S as follows. Let $i \geq 0$ be the maximum index for which $u_i = v_i$. Let C_S be the cycle obtained from S by removing u_0, u_1, \dots, u_{i-1} ($= v_0, v_1, \dots, v_{i-1}$). Then w is *to the right* of S if it is to the right of C_S when traversing C_S so that vertex u_{i+1} is visited immediately after vertex u_i . A vertex w of G is *to the left* of S if it is neither to the right of S nor a vertex of S . For the given set $B \subseteq V(G)$, let $R_B(S)$ and $L_B(S)$ be the sets of vertices in B to the right and left of S , respectively. Let $r_B(S) := |R_B(S)|$ and $\ell_B(S) := |L_B(S)|$. We drop the subscript B when




Figure 1: Two lollipops of height k . Note that the layers might have a more complicated structure than that shown here.

$B = V(G)$. By the Jordan Curve Theorem, $R_B(S)$ and $L_B(S)$ are disjoint. Note that the reverse sequence \overleftarrow{S} is also a lollipop, and $L_B(S) = R_B(\overleftarrow{S})$ and $R_B(S) = L_B(\overleftarrow{S})$.

Let S be a lollipop such that:

- (1) $r_B(S) \leq \frac{2}{3}|B|$;
- (2) subject to (1), $r_B(S)$ is maximum; and
- (3) subject to (1) and (2), $\ell(S)$ is minimum.
- (4) subject to (1), (2) and (3), $r(S)$ is maximum.

A lollipop satisfying (1) always exists, since if (r, u_1, v_1) is a face in clockwise order, then $S := (u_0 = r, u_1, v_1, v_0 = r)$ is a lollipop of height 1 with $r_B(S) = 0 \leq \frac{2}{3}|B|$.

Say S has height k . Suppose, for the sake of contradiction, that $\ell_B(S) > \frac{2}{3}|B|$. Thus $R_B(S) < \frac{1}{3}|B|$. We distinguish the following cases:

Case 1. $S = (u_0, u_1, \dots, u_{k-1}, u_k, v_k, v_{k-1}, \dots, v_1, v_0)$: Let w be the vertex that forms a face $f = (u_k, w, v_k)$ in clockwise order. By the definition of layering, $w \in V_{k+1} \cup V_k \cup V_{k-1}$.

Case 1(a). $w \in V_{k+1}$: Then $S' := (u_0, u_1, \dots, u_{k-1}, u_k, w, v_k, v_{k-1}, \dots, v_1, v_0)$ is a lollipop. Since f is a face, $R_B(S') = R_B(S)$ and $L(S') = L(S) - \{w\}$, contradicting (3).

Case 1(b). $w = u_{k-1}$: Observe that $S' := (u_0, u_1, \dots, u_{k-1}, v_k, v_{k-1}, \dots, v_1, v_0)$ is a lollipop of height $k-1$. We have $r_B(S') \leq r_B(S) + 1 < \frac{1}{3}|B| + 1 \leq \frac{2}{3}|B|$ (since $|B| \geq 3$). Hence S'

satisfies (1). If $u_k \in B$ then $r_B(S') > r_B(S)$, contradicting (2). Now assume that $u_k \notin B$. We have $\ell(S') = \ell(S)$ and $r(S') = r(S) + 1$, contradicting (4).

Case 1(c). $w = v_{k-1}$: This case is analogous to Case 1(b) except that we use $S' := (u_0, u_1, \dots, u_{k-1}, u_k, v_{k-1}, \dots, v_1, v_0)$.

Case 1(d). $w \in V_{k-1}\{u_{k-1}, v_{k-1}\}$: There is a monotone path $P = (w = z_{k-1}, z_{k-2}, \dots, z_0 = r)$ such that if some $z_i = u_i$ then $z_j = u_j$ for each $j \in [0, i]$, and if some $z_i = v_i$ then $z_j = v_j$ for each $j \in [0, i]$. Observe that both $S' := (u_0, u_1, \dots, u_{k-1}, u_k, z_{k-1}, z_{k-2}, \dots, z_1, z_0)$ and $S'' := (z_0, z_1, \dots, z_{k-1}, v_k, v_{k-1}, \dots, v_1, v_0)$ are lollipops. By assumption, $r_B(\overleftarrow{S}) = \ell_B(S) > \frac{2}{3}|B|$. We have $r_B(S') > \frac{2}{3}|B|$ since S' is a lollipop with $r_B(S') \geq r_B(S)$ and $\ell(S') < \ell(S)$. Similarly, $r_B(S'') > \frac{2}{3}|B|$. Hence $r_B(\overleftarrow{S}) + r_B(S') + r_B(S'') > 2|B|$. Thus $R_B(\overleftarrow{S}) \cap R_B(S') \cap R_B(S'') \neq \emptyset$, which is a contradiction since $R_B(S') \cap R_B(S'') \subseteq R_B(S) = L_B(\overleftarrow{S})$.

Case 1(e). $w \in V_k$: This case is analogous to Case 1(d), except that here P is a monotone path ($w = z_k, z_{k-1}, \dots, z_0 = r$), and $S' := (u_0, u_1, \dots, u_{k-1}, u_k, z_k, z_{k-1}, \dots, z_1, z_0)$ and $S'' := (z_0, z_1, \dots, z_{k-1}, z_k, v_k, v_{k-1}, \dots, v_1, v_0)$.

Case 2. $S = (u_0, u_1, \dots, u_{k-1}, u_k, u_{k+1}, v_k, v_{k-1}, \dots, v_1, v_0)$: Let w be the vertex that forms a face $f = (u_k, w, u_{k+1})$ in clockwise order. Hence $w \in V_{k+1} \cup V_k$.

Case 2(a). $w \in V_{k+1}$: This case is analogous to Case 1(a) except that $S' := (u_0, u_1, \dots, u_{k-1}, u_k, w, u_{k+1}, v_k, v_{k-1}, \dots, v_1, v_0)$.

Case 2(b). $w = v_k$: This case is analogous to Case 1(c) except with $S' := (u_0, u_1, \dots, u_{k-1}, u_k, v_k, v_{k-1}, \dots, v_1, v_0)$.

Case 2(c). $w \in V_k - \{v_k\}$: This case is analogous to Case 1(d), except that here P is a monotone path ($w = z_k, z_{k-1}, \dots, z_0$), and $S' := (u_0, u_1, \dots, u_{k-1}, u_k, z_k, z_{k-1}, \dots, z_1, z_0)$ and $S'' := (z_0, z_1, \dots, z_{k-1}, z_k, u_{k+1}, v_k, v_{k-1}, \dots, v_1, v_0)$.

Each case leads to a contradiction. Hence $r_B(S) \leq \frac{2}{3}|B|$ and $\ell_B(S) \leq \frac{2}{3}|B|$. Let G_1 be the subgraph induced by the vertices in S and to the right of S . Let G_2 be the subgraph induced by the vertices in S and to the left of S . By the Jordan Curve Theorem, no vertex to the right of S is adjacent to a vertex to the left of S . Hence $G = G_1 \cup G_2$ and (G_1, G_2) is the desired separation. \square

Lemmas 4 and 5 together prove Theorem 1 (since every planar graph with at least four vertices is a spanning subgraph of a plane triangulation).

3 Proof of Theorem 2

Theorem 2 is a special case of the following result with $H = K_5$ or $H = K_{3,3}$. A graph H is *apex* if $H - v$ is planar for some vertex v of H .

Theorem 6. *For every fixed apex graph H there is a constant $c = c(H)$ such that, for every integer k , every H -minor-free graph G is c^{k^2} -colourable such that G contains no repetitively coloured path of order at most $2k$.*

Proof. Eppstein [16] proved that for some function f (depending on H), for every H -minor-free graph G , for every vertex r of G , and for every integer $\ell \geq 0$, the set of vertices in G at distance at most ℓ from r induces a subgraph of treewidth at most $f(\ell)$. This is called the *diameter-treewidth* or *bounded local treewidth* property; also see [13, 14, 19]. Demaine and Hajiaghayi [15] strengthened Eppstein's result by showing that one can take $f(\ell) = c\ell$ for some constant $c = c(H)$.

Let G be an H -minor-free graph. By considering each connected component in turn, we may assume that G is connected. Let r be a vertex of G . Let V_0, V_1, \dots, V_p be the layering of G starting at some vertex r of G . Fix an integer $k \geq 1$. For $i \in [1, p]$, let $G_i := G[V_i \cup V_{i+1} \cup \dots \cup V_{\min\{p, i+2k-1\}}]$, and let G'_i be the minor of G obtained by contracting the connected subgraph $G[V_0 \cup V_1 \cup \dots \cup V_{i-1}]$ into a single vertex r_i . Thus G'_i is an H -minor-free graph containing G_i as a subgraph, and each vertex in G_i is at distance at most $2k$ from r_i in G'_i . By the diameter-treewidth property, G_i has treewidth at most $2ck$. By a theorem of Kündgen and Pelsmajer [29], there is a nonrepetitive 4^{2ck} -colouring ψ_i of G_i .

For each vertex v of G , define $\psi(v) := (\phi_0(v), \phi_1(v), \dots, \phi_{2k-1}(v))$, where $\phi_j(v) := \psi_i(v)$ and i is the unique integer for which $i \equiv j \pmod{2k}$ and $v \in V(G_i)$. Suppose on the contrary that G contains a repetitively coloured path $P = (v_1, \dots, v_{2t})$ of order at most $2k$ (under the colouring ψ). Thus P is contained in some G_i . Let $j := i \bmod 2k$. Hence $\psi_i(v_a) = \phi_j(v_a) = \phi_j(v_{t+a}) = \psi_i(v_{t+a})$ for each $a \in [1, t]$. That is, P is repetitively coloured by ψ_i in the colouring of G_i . This contradiction proves that G contains no repetitively coloured path under ψ . The number of colours is $(4^{2ck})^{2k} = (4^{4c})^{k^2}$. \square

Graphs embeddable on a fixed surface exclude a fixed apex graph as a minor [16]. Thus Theorem 6 implies:

Corollary 7. *For every fixed surface Σ there is a constant $c = c(\Sigma)$ such that, for every integer $k \geq 1$, every graph G embeddable in Σ is c^{k^2} -colourable such that G contains no repetitively coloured path of order at most $2k$.*

4 Open Problems

Our research suggests two open problems:

1. Is $\pi(G) \in o(\log n)$ for every planar graph G with n vertices?
2. Is there a polynomial function f such that for every integer $k \geq 1$ every planar graph G is $f(k)$ -colourable such that G contains no repetitively coloured path of order at most $2k$?

Finally, we mention a class of planar graphs that seem difficult to nonrepetitively colour. Let T be a tree rooted at a vertex r . Let V_i be the set of vertices in T at distance i from r . Draw T in the plane with no crossings. Add a cycle on each V_i in the cyclic order defined by the drawing to create a planar graph G_T . It is open whether $\pi(G_T) \leq c$ for some constant c independent of T . Note that this class of planar graphs includes examples with unbounded degree and unbounded treewidth.

References

- [1] MICHAEL O. ALBERTSON, GLENN G. CHAPPELL, HAL A. KIERSTEAD, ANDRÉ KÜNDGEN, AND RADHIKA RAMAMURTHI. Coloring with no 2-colored P_4 's. *Electron. J. Combin.*, 11 #R26, 2004. http://www.combinatorics.org/Volume_11/Abstracts/v11i1r26.html. MR: 2056078.
- [2] NOGA ALON AND JAROSŁAW GRYTCZUK. Breaking the rhythm on graphs. *Discrete Math.*, 308:1375–1380, 2008. doi: [10.1016/j.disc.2007.07.063](https://doi.org/10.1016/j.disc.2007.07.063). MR: 2392054.
- [3] NOGA ALON, JAROSŁAW GRYTCZUK, MARIUSZ HAŁUSZCZAK, AND OLIVER RIORDAN. Nonrepetitive colorings of graphs. *Random Structures Algorithms*, 21(3-4):336–346, 2002. doi: [10.1002/rsa.10057](https://doi.org/10.1002/rsa.10057). MR: 1945373.
- [4] JÁNOS BARÁT AND PÉTER P. VARJÚ. On square-free vertex colorings of graphs. *Studia Sci. Math. Hungar.*, 44(3):411–422, 2007. doi: [10.1556/SScMath.2007.1029](https://doi.org/10.1556/SScMath.2007.1029). MR: 2361685.
- [5] JÁNOS BARÁT AND PÉTER P. VARJÚ. On square-free edge colorings of graphs. *Ars Combin.*, 87:377–383, 2008. MR: 2414029.
- [6] JÁNOS BARÁT AND DAVID R. WOOD. Notes on nonrepetitive graph colouring. *Electron. J. Combin.*, 15:R99, 2008. http://www.combinatorics.org/Volume_15/Abstracts/v15i1r99.html. MR: 2426162.

- [7] BOŠTJAN BREŠAR, JAROSŁAW GRYTCZUK, SANDI KLAVŽAR, STANISŁAW NIWCZYK, AND IZTOK PETERIN. Nonrepetitive colorings of trees. *Discrete Math.*, 307(2):163–172, 2007. doi: [10.1016/j.disc.2006.06.017](https://doi.org/10.1016/j.disc.2006.06.017). MR: [2285186](#).
- [8] BOŠTJAN BREŠAR AND SANDI KLAVŽAR. Square-free colorings of graphs. *Ars Combin.*, 70:3–13, 2004. MR: [2023057](#).
- [9] PANAGIOTIS CHEILARIS, ERNST SPECKER, AND STATHIS ZACHOS. Neochromatic. *Comment. Math. Univ. Carolin.*, 51(3):469–480, 2010. <http://www.dml.cz/dmlcz/140723>. MR: [2741880](#).
- [10] JAMES D. CURRIE. There are ternary circular square-free words of length n for $n \geq 18$. *Electron. J. Combin.*, 9(1), 2002. http://www.combinatorics.org/Volume_9/Abstracts/v9i1n10.html. MR: [1936865](#).
- [11] JAMES D. CURRIE. Pattern avoidance: themes and variations. *Theoret. Comput. Sci.*, 339(1):7–18, 2005. doi: [10.1016/j.tcs.2005.01.004](https://doi.org/10.1016/j.tcs.2005.01.004). MR: [2142070](#).
- [12] SEBASTIAN CZERWIŃSKI AND JAROSŁAW GRYTCZUK. Nonrepetitive colorings of graphs. 6th Czech-Slovak International Symposium on Combinatorics, Graph Theory, Algorithms and Applications. *Electron. Notes Discrete Math.*, 28:453–459, 2007. doi: [10.1016/j.endm.2007.01.063](https://doi.org/10.1016/j.endm.2007.01.063). MR: [2324051](#).
- [13] ERIK D. DEMAIN, FEDOR V. FOMIN, MOHAMMAD TAGHI HAJIAGHAYI, AND DIMITRIOS M. THILIKOS. Bidimensional parameters and local treewidth. *SIAM J. Discrete Math.*, 18(3):501–511, 2004/05. doi: [10.1137/S0895480103433410](https://doi.org/10.1137/S0895480103433410). MR: [2134412](#).
- [14] ERIK D. DEMAIN AND MOHAMMAD TAGHI HAJIAGHAYI. Diameter and treewidth in minor-closed graph families, revisited. *Algorithmica*, 40(3):211–215, 2004. doi: [10.1007/s00453-004-1106-1](https://doi.org/10.1007/s00453-004-1106-1). MR: [2080518](#).
- [15] ERIK D. DEMAIN AND MOHAMMAD TAGHI HAJIAGHAYI. Equivalence of local treewidth and linear local treewidth and its algorithmic applications. In *Proc. 15th Annual ACM-SIAM Symposium on Discrete Algorithms* (SODA '04), pp. 840–849. SIAM, 2004. <http://dl.acm.org/citation.cfm?id=982792.982919>.
- [16] DAVID EPPSTEIN. Diameter and treewidth in minor-closed graph families. *Algorithmica*, 27(3-4):275–291, 2000. doi: [10.1007/s004530010020](https://doi.org/10.1007/s004530010020). MR: [1759751](#).
- [17] GUILLAUME FERTIN, ANDRÉ RASPAUD, AND BRUCE REED. Star coloring of graphs. *J. Graph Theory*, 47(3):163–182, 2004. doi: [10.1002/jgt.20029](https://doi.org/10.1002/jgt.20029). MR: [2089462](#).
- [18] FRANCESCA FIORENZI, PASCAL OCHEM, PATRICE OSSONA DE MENDEZ, AND XUDING ZHU. Thue choosability of trees. *Discrete Applied Math.*, 159(17):2045–2049, 2011. doi: [10.1016/j.dam.2011.07.017](https://doi.org/10.1016/j.dam.2011.07.017). MR: [2832329](#).

- [19] MARTIN GROHE. Local tree-width, excluded minors, and approximation algorithms. *Combinatorica*, 23(4):613–632, 2003. doi: [10.1007/s00493-003-0037-9](https://doi.org/10.1007/s00493-003-0037-9). MR: 2046826.
- [20] JAROSŁAW GRYTCZUK. Thue-like sequences and rainbow arithmetic progressions. *Electron. J. Combin.*, 9(1):R44, 2002. http://www.combinatorics.org/Volume_9/Abstracts/v9i1r44.html. MR: 1946146.
- [21] JAROSŁAW GRYTCZUK. Nonrepetitive colorings of graphs—a survey. *Int. J. Math. Math. Sci.*, 74639, 2007. doi: [10.1155/2007/74639](https://doi.org/10.1155/2007/74639). MR: 2272338.
- [22] JAROSŁAW GRYTCZUK. Nonrepetitive graph coloring. In *Graph Theory in Paris*, Trends in Mathematics, pp. 209–218. Birkhauser, 2007.
- [23] JAROSŁAW GRYTCZUK. Thue type problems for graphs, points, and numbers. *Discrete Math.*, 308(19):4419–4429, 2008. doi: [10.1016/j.disc.2007.08.039](https://doi.org/10.1016/j.disc.2007.08.039). MR: 2433769.
- [24] JAROSŁAW GRYTCZUK, JAKUB KOZIK, AND PIOTR MICEK. A new approach to non-repetitive sequences. 2011. arXiv: [1103.3809](https://arxiv.org/abs/1103.3809). To appear in *Random Structures Algorithms*.
- [25] JAROSŁAW GRYTCZUK, JAKUB PRZYBYŁO, AND XUDING ZHU. Nonrepetitive list colourings of paths. *Random Structures Algorithms*, 38(1-2):162–173, 2011. doi: [10.1002/rsa.20347](https://doi.org/10.1002/rsa.20347). MR: 2768888.
- [26] JOCHEN HARANTA AND STANISLAV JENDROL'. Nonrepetitive vertex colorings of graphs. *Discrete Math.*, 312(2):374–380, 2012. doi: [10.1016/j.disc.2011.09.027](https://doi.org/10.1016/j.disc.2011.09.027).
- [27] FRÉDÉRIC HAVET, STANISLAV JENDROL', ROMAN SOTÁK, AND ERIKA ŠKRABUL'ÁKOVA. Facial non-repetitive edge-coloring of plane graphs. *J. Graph Theory*, 66(1):38–48, 2011. doi: [10.1002/jgt.20488](https://doi.org/10.1002/jgt.20488). MR: 2742187.
- [28] STANISLAV JENDROL AND ERIKA ŠKRABUL'ÁKOVÁ. Facial non-repetitive edge colouring of semiregular polyhedra. *Acta Univ. M. Belii Ser. Math.*, 15:37–52, 2009. <http://actamath.savbb.sk/acta1503.shtml>. MR: 2589669.
- [29] ANDRE KÜNDGEN AND MICHAEL J. PELSMAJER. Nonrepetitive colorings of graphs of bounded tree-width. *Discrete Math.*, 308(19):4473–4478, 2008. doi: [10.1016/j.disc.2007.08.043](https://doi.org/10.1016/j.disc.2007.08.043). MR: 2433774.
- [30] FEDOR MANIN. The complexity of nonrepetitive edge coloring of graphs, 2007. arXiv: [0709.4497](https://arxiv.org/abs/0709.4497).
- [31] DÁNIEL MARX AND MARCUS SCHAEFER. The complexity of nonrepetitive coloring. *Discrete Appl. Math.*, 157(1):13–18, 2009. doi: [10.1016/j.dam.2008.04.015](https://doi.org/10.1016/j.dam.2008.04.015). MR: 2479374.

- [32] JAROSLAV NEŠETŘIL AND PATRICE OSSONA DE MENDEZ. Colorings and homomorphisms of minor closed classes. In BORIS ARONOV, SAUGATA BASU, JÁNOS PACH, AND MICHA SHARIR, eds., *Discrete and Computational Geometry, The Goodman-Pollack Festschrift*, vol. 25 of *Algorithms and Combinatorics*, pp. 651–664. Springer, 2003. MR: [2038495](#).
- [33] JAROSLAV NEŠETŘIL, PATRICE OSSONA DE MENDEZ, AND DAVID R. WOOD. Characterisations and examples of graph classes with bounded expansion. *European J. Combinatorics*, 33(3):350–373, 2011. doi: [10.1016/j.ejc.2011.09.008](https://doi.org/10.1016/j.ejc.2011.09.008).
- [34] WESLEY PEGDEN. Highly nonrepetitive sequences: winning strategies from the local lemma. *Random Structures Algorithms*, 38(1-2):140–161, 2011. doi: [10.1002/rsa.20354](https://doi.org/10.1002/rsa.20354). MR: [2768887](#)
- [35] ANDRZEJ PEZARSKI AND MICHAŁ ZMARZ. Non-repetitive 3-coloring of subdivided graphs. *Electron. J. Combin.*, 16(1):N15, 2009. http://www.combinatorics.org/Volume_16/Abstracts/v16i1n15.html. MR: [2515755](#).
- [36] AXEL THUE. Über unendliche Zeichenreihen. *Norske Vid. Selsk. Skr. I. Mat. Nat. Kl. Christiania*, 7:1–22, 1906.
- [37] DAVID R. WOOD. Acyclic, star and oriented colourings of graph subdivisions. *Discrete Math. Theor. Comput. Sci.*, 7(1):37–50, 2005. <http://www.dmtcs.org/dmtcs-ojs/index.php/dmtcs/article/view/60>. MR: [2164057](#).